

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/323918707>

Body fat in prepubertal boys: different training program`s designs

Poster · March 2018

CITATIONS

0

READS

56

6 authors, including:



Ana Ruivo Alves

Polytechnic Institute of Beja

22 PUBLICATIONS 20 CITATIONS

[SEE PROFILE](#)



Carlos Marta

Polytechnic Institute of Guarda

37 PUBLICATIONS 147 CITATIONS

[SEE PROFILE](#)



Henrique P. Neiva

Universidade da Beira Interior

85 PUBLICATIONS 273 CITATIONS

[SEE PROFILE](#)



Júlío Martins

Universidade da Beira Interior

122 PUBLICATIONS 105 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Biomechanic in strength training [View project](#)



Pacing and Soccer Training [View project](#)

BODY FAT IN PREPUBERTAL BOYS: DIFFERENT TRAINING PROGRAM'S DESIGNS

ANA R. ALVES¹, CARLOS MARTA^{3,4}, HENRIQUE P. NEIVA^{1,2}, DANIEL A. MARINHO^{1,2}, JÚLIO MARTINS ^{1,2}, AND MÁRIO C. MARQUES^{1,2}

¹Department of Sport Sciences, University of Beira Interior, Covilhã, Portugal; ²Research Center in Sport Sciences, Health Sciences and Human Development, CIDESD, Covilhã, Portugal; ³Department of Sport Sciences, Guarda Polytechnique Institute, Guarda, Portugal; ⁴Research Unit for Inland Development, UDI, Guarda, Portugal. —

INTRODUCTION

Research studies agreed that strength and aerobic training either combined or performed separately promote bone density, aerobic capacity, and explosive strength improvements in childhood. Nevertheless, there still is uncertainty regarding the best training program to be implemented that efficiently improve body fat percentage. The current study aimed to compare different training program's designs on body fat percentage in prepubertal boys.

METHODS

One hundred twenty-three boys (10.93 ± 0.57 years) were randomly assigned into five experimental groups to perform different training protocols for 8 weeks: strength-only (SG), aerobic-only (AG), inter-session concurrent aerobic-strength training (ASG_2), intra-session concurrent aerobic-strength training (ASG), intra-session concurrent strength-aerobic training (SAG), and a control group, no training (CG). SG, AG, ASG and SAG groups performed single sessions two days per week. ASG_2 group performed on different days each week (two and two days per week). The strength training program comprised plyometric exercises (medicine ball throws and jumps) and sprint running. The aerobic training program comprised the 20m shuttle run exercise. Body fat percentage was assessed using a body composition analyzer (Tanita TBF-300A) before and after the 8-weeks of the training program.

RESULTS

Body fat percentage showed significant decrements from pre- to the post-training in the ASG_2 (17.6%, $p=0.004$), SG (16.1%, $p=0.000$), and SAG (17.7%, $p=0.000$) groups. There was an impairment in the ASG (4.2%, $p=0.492$) and control group CG (16.8%, $p=0.000$). No differences were presented in the AG ($p=0.053$).

CONCLUSIONS

The order of intra-session concurrent training influenced body fat percentage changes. Moreover, performing intra-session concurrent strength and aerobic training is more useful than strength or aerobic training only and concurrent training in different sessions to decrease body fat percentage in prepubertal boys. These results have a meaningful interest to optimized school-based fat loss exercise programs in childhood.

ACKNOWLEDGMENTS

This project was supported by the National Funds through FCT - Portuguese Foundation for Science and Technology (UID/DTP/04045/2013) - and the European Fund for regional development (FEDER) allocated by European Union through the COMPETE 2020 Programme (POCI-01-0145-FEDER-006969), and it was supported by the Project NanoSTIMA: Macro-to-Nano Human Sensing, Towards Integrated Multimodal Health Monitoring and Analytics, NORTE-01-0145-FEDER-000016, co-financed by European Fund for Regional Development (FEDER) - NORTE 2020